



Vol. XV &amp; Issue No. 11 November - 2022

INDUSTRIAL ENGINEERING JOURNAL

## AN EFFICIENT SUPPLIER SELECTION FOR INDIAN PHARMACEUTICAL INDUSTRY BY APPLYING FUZZY TOPSIS METHOD

**Dashrath Kumar Patidar**

**Dr. V. K. Sukhwani**

**Dr. A. C. Shukla**

### Abstract

*Supplier selection is one of the critical processes in supply chain management which is associated with the flow of goods and services from the supplier of raw material to the last consumer. It is a multi-criteria problem which includes both qualitative and quantitative factors (criteria). Under many conditions the values of the criteria are often inaccurately defined for the decision makers. To reduce this problem, fuzzy multi-criteria decision making approach can be used. In this paper, a multi-criteria decision-making approach has been used for selecting supplier under uncertainty. The aim of this study is to develop a method to evaluate suppliers in supply chain based on technique for order preference by similarity to ideal solution method (TOPSIS) to select the most efficient supplier. This paper presents a successful application of FUZZY TOPSIS to an actual supplier selection problem in Indian pharmaceutical industry.*

**Keywords:** supplier selection, supply chain management, multi-criteria decision-making, FUZZY TOPSIS method

### INTRODUCTION

Suppliers are key aspect of modern supply chain and play important role in the success or failure of any business. Supplier has substantial impact on the efficiency and effectiveness of the entire supply chain. Therefore, selecting the efficient suppliers can result in reduced cost, decreased supplying risk and improved quality. In the present market of global competition, it has become necessary for the manufactures to reduce their fixed costs and supply the goods at the right place in the right time to sustain the competition and survive in the market. India supplies 20% of global generic medicines market export in terms of volume, making the country the largest provider of generic medicines globally and expected to expand even further in coming years. It is likely that the manufacturer allocates more than 70% of its total production on purchased services and materials (Athena Forghani, 2018). It would not be easy for the decision makers to select best suppliers who can fulfill all the needs of the firm based on different criteria.

The pharmacies are ready to utilize a tremendous amount of pharmacy's financial resources in supplier selection process. In return, the pharmacy managers expect the reliable supplier in all the way to fulfill the service, on-time delivery at low-cost compared to other suppliers (P. Kelle, J. Woosley, and H. Schneider 2012). Other point to be considered is that, as a multiple criteria decision-making (MCDM) problem, the supplier selection would lie under the effect of many qualitative and quantitative conflicting factors. In general, multi criteria group decision-making (MCDM) problems are frequently used in this type of cases.

This study intends to adopt a qualitative method by using fuzzy TOPSIS algorithm with triangular fuzzy number. The

extent method has been used to calculate the weight of criteria and Fuzzy TOPSIS method has been used to evaluate rate of alternatives. Fuzzy TOPSIS is a method that can help in objective and systematic evaluation of alternatives on multiple criteria. The fuzzy TOPSIS method hinges on the concept that the chosen alternative should have the shortest distance from the positive ideal solution (PIS) and the farthest from the negative ideal solution (NIS). The Fuzzy TOPSIS approach has been applied to various MCDM problems like location selection (Farimah mokhatab rafiei, 2012), optimal location airport fire stations (Mehmet Sevkli, 2014).

### LITERATURE REVIEW

According to Tahiri et al.2008, supplier selection problem has become one of the most important issues for establishing an effective supply chain system". Indeed, supplier selection and evaluation represents one of the significant roles of purchasing and supply management function (Kumara et al. 2003) contend that strategic partnership with the right supplier must be integrated within the supply chain to contain costs, improve quality and flexibility to meet end-customers' value and reduce lead time at different stages of the supply chain (Chris I., Bell-Hanyes, 2010).

The analysis of criteria for selection and measuring the performance of suppliers has been the focus of many academicians and purchasing practitioners since 1960s. (Weber,Current, Benton, 1991) Based on (Dickson's 1966) empirical study, 23 criteria were identified which purchasing managers generally consider when selecting a supplier. Of the identified criteria, quality, on-time delivery, and supplier's performance history were found vital in supplier selection regardless of the type of purchasing environment.

A two-stage model was developed by ( Liao and Kao, 2011) by applying Fuzzy TOPSIS and multi-choice GP for selection the appropriate suppliers and allocating the orders. Also Fuzzy TOPSIS and multi-choice GP were used (Rouyendegh and Saputro, 2014) in a fertilizer and chemical producing company.

The Fuzzy TOPSIS approach has been applied to various MCDM problems ranging from facility location selection (T.C. Chu, 2002), robot selection (T.C. Chu, and Y.C. Lin, 2003), selection of system analysis engineer for a software company and choosing optimal initial training aircraft in Air Force Academy in Taiwan (T.C. Wang, and T.H. Chang, 2007), to service quality in hotel industry (J.M. Benitez, J.C. Martin, and C. Roman, 2007), plant layout design problem (T. Yang and C.C. Hung, 2007), transshipment site selection (S. Onut, and S. Sonar, 2008), and machine tool selection problem (M. Yurdakul, and Y.T. Ic, 2008).

The AHP approach was used to carry out an analysis of strategic supplier selection and evaluation in a generic pharmaceutical firm supply chain (Enyinda, et al., 2010). The researchers developed a model to aid them in the evaluation and selection of the important criteria and hence the best supplier for a pharmaceutical manufacturing firm. The selected criteria for the evaluation were regulatory compliance, quality, cost, service, supplier profile and risk. The researchers recommended that supplier selection process and evaluation represents one of the key activities that organizations must integrate into their core strategic decisions. Based on their research findings, the regulatory compliance selection criterion was most favored, followed by quality, risk, cost, supplier profile, and service. The model also enabled the researchers to select the best supplier for the case company (David Asamoah, J. A. 2012).

## RESEARCH METHODOLOGY

The study approach comprises of two steps. In the first step, various criteria for suppliers rating selection are identified. In second step the experts in pharmaceutical industry were asked to give linguistic ratings to the criteria and the alternatives. The major supplier selection criteria in pharmaceutical companies are grouped under sixteen factors and a questionnaire was prepared to collect information about the importance of the criteria. Then the Fuzzy TOPSIS method has been applied which here consist of supplier's criteria as given below.

**Supplier's criteria:** In pharmaceutical companies, the main criterions are grouped under sixteen factors, which are shown in table-1 (Athena Forghani, 2018). Furthermore, an online questionnaire was prepared to collect rating information of selected criteria assessment and alternative assessment under study. Rating information of alternative assessment consists of a scale from 1 to 5, where 1 shows very low, 2 shows low, 3 shows medium, 4 shows high and 5 shows very high. Whereas, rating information of criteria assessment consists of a scale from 1 to 5, where 1 shows very poor, 2 shows poor, 3 shows fair, 4 shows good and 5 shows very good. Then the Fuzzy TOPSIS method was applied to get the alternative rating and to find the proper weightage to various criteria.

**Table 1: Factors for supplier selection**

Sr. no.	Factors	Significance
1	Response time ( $C_1$ )	Commitment of supplier towards fastest response time.
2	On time delivery ( $C_2$ )	Increases awareness about on time delivery importance.
3	Product quality ( $C_3$ )	Commitment by supplier towards product quality.
4	Product price ( $C_4$ )	Supplier pricing strategies could shape our overall profitability for the future.
5	Complete shipping document ( $C_5$ )	Properly completed documentation will help your shipment reach its destination on time.
6	Reputation ( $C_6$ )	A supplier good reputation is definitely preferred for better business.
7	Past record documentation ( $C_7$ )	Keeping clear past record documentation for just good relations.
8	Geographic location ( $C_8$ )	Global it is important to consider the geographical location of the supply.
9	Social factor ( $C_9$ )	Social factor represents another important set of influences on consumer behavior.
10	Environment factor ( $C_{10}$ )	Environment respective image in the view of society, on any supplier.
11	Service warranty ( $C_{11}$ )	To better understand the importance of product warranties in supplier today.
12	Flexibility (response to change) ( $C_{12}$ )	Supplier flexibility refers to the ability of the supplier organization to respond to the changing demand.
13	Overall personal capability ( $C_{13}$ )	Better overall personal capability of any supplier.
14	Supplier profile ( $C_{14}$ )	Supplier profile important in business for market.
15	Financial status ( $C_{15}$ )	Commitment of supplier to take financial rise.
16	Communication system ( $C_{16}$ )	Effective communication between buyers and suppliers helps support long terms goals.

**Table 2: Detailed profile of the criteria ratings of decision makers.**

DM	Experts	Designation	Department/Area of expertise	Experience (years)
DM <sub>1</sub>	1	Head of quality control	Quality	14
DM <sub>2</sub>	2	Medical representative	Sales	9
DM <sub>3</sub>	3	Procurement manager	Purchase	19
DM <sub>4</sub>	4	Purchase executive	Purchase	8

**The FUZZY TOPSIS theory details**


The *Technique for order of preference by similarity to ideal solution* (TOPSIS) is a multi criteria decision analysis method which was originally developed by Hwang and Yoon in 1981 with further developments by Yoon in 1987 and Hwang, Lai and Liu in 1993 (Tonekaboni, 2012). The technique called Fuzzy TOPSIS (Technique for Order Preference by Similarity to Ideal Situation) can be used to evaluate multiple alternatives against the selected criteria. In the TOPSIS approach an alternative that is nearest to the Fuzzy Positive Ideal Solution (FPIS) and farthest from the Fuzzy Negative Ideal Solution (FNIS) is chosen as optimal. An FPIS is composed of the best performance values for each alternative whereas the FNIS consists of the worst performance values (Tonekaboni, 2012). The definitions of this method are as follow:

**Definition 1:** A fuzzy set  $\tilde{a}$  in a universe of discourse X is characterized by a membership function  $\mu_{\tilde{a}}(x)$  that maps each element x in X to a real number in the interval [0,1]. The function value  $\mu_{\tilde{a}}(x)$  is termed the grade of membership of x in  $\tilde{a}$ . The nearer the value of  $\mu_{\tilde{a}}(x)$  to unity, the higher the grade of membership of x in  $\tilde{a}$  (Kore, 2017).

**Definition 2:** A triangular fuzzy number is represented as a triplet  $\tilde{a} = (a_1, a_2, a_3)$ . The membership function  $\mu_{\tilde{a}}(x)$  of triangular fuzzy number  $\tilde{a}$  (Kore, 2017).

A fuzzy set  $\tilde{a}$ , membership function  $\mu_{\tilde{a}}$  that maps each element x in X to a real number in the interval [0, 1]. A triangular fuzzy number is represented  $\tilde{a} = (a_1, a_2, a_3)$ .

**Figure 1:** Triangular Fuzzy number system, (Kore, 2017)



$a_2$  gives the maximal grade of  $\mu_a$  that  $\mu_a=1$

$a_1$  gives the minimal grade of  $\mu_a$  that  $\mu_a=0$

$a_1$  and  $a_3$  are the lower and upper bounds of the available area for the evaluation data.

$$\mu_{\tilde{a}}(x) = \begin{cases} (x-a_1) / (a_2 - a_1) & \text{if } a_1 \leq x \leq a_2 \\ (a_3 - x) / (a_3 - a_2) & \text{if } a_2 \leq x \leq a_3 \\ 0 & \text{otherwise} \end{cases}$$

$$\mu_{\tilde{a}}(x) = \begin{cases} 0 & \text{if } x < a_1 \\ (x - a_1) / (a_2 - a_1) & \text{if } a_2 \geq x \geq a_1 \\ (x - a_2) / (a_3 - a_2) & \text{if } a_2 \geq x \geq a_3 \\ 0 & \text{if } x > a_3 \end{cases}$$

The distance between fuzzy triangular numbers

Let  $\tilde{a} = (a_1, a_2, a_3)$  and  $b = (b_1, b_2, b_3)$  be two triangular Fuzzy numbers. The distance between them is given using the vertex method by:

$$d(\tilde{a}, \tilde{b}) = \sqrt{1/3[(a_1-b_1)^2+(a_2-b_2)^2+(a_3-b_3)^2]} \dots \dots \dots (1)$$

The various steps of fuzzy TOPSIS are presented as follows:

**Step 1:** assignment of rating to the criteria and the alternatives.

Let us assume there are J possible candidates called  $S = \{S_1, S_2, \dots, S_j\}$  which are to be evaluated against n criteria  $C = \{C_1, C_2, \dots, C_n\}$ . The criteria weights are denoted by  $W_i (i=1, \dots, m)$ . The performance ratings of each decision maker  $D_k (k=1, \dots, K)$  for each alternative  $A_j (j=1, \dots, n)$  with respect to criteria  $C_i (i=1, \dots, m)$  are denoted by  $\tilde{R}_k = \tilde{x}_{ijk} = 1, \dots, m, j = 1, \dots, n, k = 1, \dots, k$  with membership function  $\mu_{Rk}$ .

**Step 2:** Compute aggregate fuzzy ratings for the criteria and the alternatives.

If the fuzzy ratings of all decision makers is described as triangular fuzzy number  $\tilde{R}_k = (a_k, b_k, c_k), k=1, \dots, K$ , then the aggregated fuzzy rating is given by  $\tilde{R} = (a, b, c), k = 1, \dots, K$ , where  $a = \min_k \{a_k\}, b = \frac{1}{K}, c = \max_k \{c_k\} \dots \dots \dots (2)$

If the fuzzy rating and importance weight of the  $k^{th}$  decision maker are  $\tilde{x}_{jk} = (a_{ijk}, b_{ijk}, c_{ijk})$  and  $\tilde{w}_{ijk} = (w_{jk1}, w_{jk2}, w_{jk3}), i=1, 2, \dots, m; j=1, 2, \dots, n$  respectively, then the aggregated fuzzy ratings ( $\tilde{X}_{ij}$ ) of alternatives with respect to each criteria are given by  $\tilde{x}_{ij} = (a_{ij}, b_{ij}, c_{ij})$  where

$$a_{ij} = \min_k \{a_{ijk}\}, b_{ij} = \frac{1}{K} \sum_{k=1}^K \{b_{ijk}\}, c_{ij} = \max_k \{c_{ijk}\} \dots \dots \dots (3)$$

The aggregated fuzzy weights ( $\tilde{w}_{ij}$ ) of each criterion are calculated as  $\tilde{w}_j = (w_{j1}, w_{j2}, w_{j3})$  where:

$$W_{j1} = \min_k \{w_{jk1}\}, w_{j2} = \frac{1}{K} \sum_{k=1}^K \{w_{jk2}\}, w_{j3} = \max_k \{w_{jk3}\} \dots \dots \dots (4)$$

**Step 3:** Compute the fuzzy decision matrix: The fuzzy decision matrix for the alternatives ( $\tilde{D}$ ) and the criteria ( $\tilde{W}$ ) is constructed as follows:

$$\tilde{D} = \begin{matrix} & C_1 & C_2 & \dots & C_{16} \\ S_1 & x_{11} & x_{12} & \dots & x_{1n} \\ S_2 & x_{21} & x_{22} & \dots & x_{2n} \\ \dots & \dots & \dots & \dots & \dots \\ S_n & x_{n1} & x_{n2} & \dots & x_{nn} \end{matrix} \quad I = 1, \dots, m; j = 1, \dots, n$$

$$\tilde{w} = (\tilde{w}_1, \tilde{w}_2, \dots, \tilde{w}_n)$$

**Step 4:** Normalize the fuzzy decision matrix: The raw data are normalized using linear scale transformation to bring the

various criteria scales into a comparable scale. The normalized fuzzy decision matrix  $\tilde{R}$  is given by:

$$\tilde{R} = [\tilde{r}_{ij}]_{m \times n}, i = 1, \dots, m, j = 1, \dots, n$$

Where,

$$\tilde{r}_{ij} = (a_{ij}/c_j^*, b_{ij}/c_j^*, c_{ij}/c_j^*) \text{ and } c_j^* = \max_i c_{ij}$$

(benefit criteria).....(5)

$$\tilde{r}_{ij} = (\bar{a}_{ij}/c_{ij}, \bar{a}_{ij}/b_{ij}, \bar{a}_{ij}/a_{ij}) \text{ and } \bar{a}_{ij} = \min_i a_{ij}$$

(cost criteria).....(6)

**Step 5:** Compute the weighted normalized matrix: The weighted normalized matrix  $\tilde{P}$  for criteria is computed by multiplying the weights ( $\tilde{w}_j$ ) of evaluation criteria with the normalized fuzzy decision matrix  $\tilde{r}_{ij}$

$$\tilde{P} = [p_{ij}]_{m \times n}, i = 1, \dots, m, j = 1, \dots, n$$

Where  $\tilde{p}_{ij} = \tilde{r}_{ij}(\cdot)\tilde{w}_j$ .....(7)

**Step 6:** Compute the fuzzy positive ideal solution (FPIS) and fuzzy negative ideal solution (FNIS): The FPIS and FNIS of the alternatives are computed as follows:

$$A^+ = (\tilde{p}_1^+, \tilde{p}_2^+, \dots, \tilde{p}_n^+), \text{ where } \tilde{p}_j^+ = \max_i(p_{ij}), i = 1, \dots, m, j = 1, \dots, n.....(8)$$

$$A^- = (\tilde{p}_1^-, \tilde{p}_2^-, \dots, \tilde{p}_n^-), \text{ where } \tilde{p}_j^- = \min_i(p_{ij}), i = 1, \dots, m, j = 1, \dots, n.....(9)$$

**Step 7:** Compute the distance of alternative from FPIS and FNIS: The distance ( $d_i^+, d_i^-$ ) of each weighted alternative  $I = 1, \dots, m$  from the FPIS and the FNIS is computed as follows:

$$d_i^+ = \sum_{j=1}^n d_p(\tilde{p}_{ij}, \tilde{p}_j^+), i = 1, \dots, m.....(10)$$

$$d_i^- = \sum_{j=1}^n d_p(\tilde{p}_{ij}, \tilde{p}_j^-), i = 1, \dots, m.....(11)$$

Where,  $d_p(\tilde{a}, \tilde{b})$  is the distance measurement between two fuzzy number  $\tilde{a}$  and  $\tilde{b}$

**Step 8:** Compute the closeness coefficient ( $CC_i$ ) of each alternative: The closeness coefficient  $CC_i$  represents the distances to the fuzzy positive ideal solution ( $A^+$ ) and the fuzzy negative ideal solution ( $A^-$ ) simultaneously. The closeness coefficient of each alternative is calculated as:

$$CC_i = d_i^- / (d_i^+ + d_i^-), I = 1, 2, \dots, m.....(12)$$

**THE CASE STUDY**

Fuzzy set theory was firstly introduced by Zadeh 1965 (Vinod Yadav, M. K., 2016). Conversion scales are applied to transform the linguistic terms into fuzzy numbers. Usually apply a scale of 1 to 9 for rating the criteria and the alternatives. The intervals are chosen so as to have a uniform representation from 1 to 9 for the fuzzy triangular numbers used for the five linguistic ratings.

**Table 3: Fuzzy ratings for linguistic variables**

Fuzzy number	Criteria assessment	Alternative assessment
(1,1,3)	Very Poor (VP)	Very low (VL)

(1,3,5)	Poor (P)	Low (L)
(3,5,7)	Fair (F)	Medium (M)
(5,7,9)	Good (G)	High (H)
(7,9,9)	Very Good (VG)	Very high (VH)

**Steps to illustrate Fuzzy TOPSIS method**

Let's say, the decision group has K members and the  $i^{th}$  alternative on  $j^{th}$  criterion. The fuzzy rating and importance weight of the  $k^{th}$  decision maker, about the  $i^{th}$  alternative on  $j^{th}$  criterion.

**Step 1:** We have five alternatives such as  $S_1, S_2, S_3, S_4$  and  $S_5$  for comparison with sixteen criteria such as  $C_1, C_2, C_3, \dots, C_{16}$  also we have four decision makers namely  $DM_1, DM_2, DM_3$  and  $DM_4$ .

**Step 2:** Criteria rating by decision makers.

**Table 4: Criteria rating**

Criteria	DM <sub>1</sub>	DM <sub>2</sub>	DM <sub>3</sub>	DM <sub>4</sub>
C <sub>1</sub>	G	VG	VG	G
C <sub>2</sub>	VG	VG	G	VG
C <sub>3</sub>	VG	VG	G	VG
C <sub>4</sub>	G	P	F	F
C <sub>5</sub>	G	VG	VG	VG
C <sub>6</sub>	F	VG	F	F
C <sub>7</sub>	F	VP	F	P
C <sub>8</sub>	P	P	P	VP
C <sub>9</sub>	G	G	F	F
C <sub>10</sub>	VG	VG	F	G
C <sub>11</sub>	G	G	G	G
C <sub>12</sub>	F	F	G	F
C <sub>13</sub>	G	P	F	F
C <sub>14</sub>	G	G	F	G
C <sub>15</sub>	F	G	F	F
C <sub>16</sub>	G	VG	G	VG

**Step 3:** Now, decision makers are rating the alternatives as shown in table-5.

Table 5: Alternative rating

Criteria ↓	S <sub>1</sub>				S <sub>2</sub>				S <sub>3</sub>				S <sub>4</sub>				S <sub>5</sub>			
	DM <sub>1</sub>	DM <sub>2</sub>	DM <sub>3</sub>	DM <sub>4</sub>	DM <sub>1</sub>	DM <sub>2</sub>	DM <sub>3</sub>	DM <sub>4</sub>	DM <sub>1</sub>	DM <sub>2</sub>	DM <sub>3</sub>	DM <sub>4</sub>	DM <sub>1</sub>	DM <sub>2</sub>	DM <sub>3</sub>	DM <sub>4</sub>	DM <sub>1</sub>	DM <sub>2</sub>	DM <sub>3</sub>	DM <sub>4</sub>
C <sub>1</sub>	M	VH	VH	L	H	M	H	L	H	H	H	VH	H	VH	M	H	L	L	L	M
C <sub>2</sub>	H	VH	H	L	H	VH	H	L	H	H	VH	H	H	H	M	H	H	M	M	M
C <sub>3</sub>	M	VH	M	H	H	VH	VH	H	M	M	VH	H	H	H	H	H	H	L	H	VH
C <sub>4</sub>	VL	VL	H	H	M	L	VH	H	M	H	VH	H	M	M	H	VH	M	VH	L	VH
C <sub>5</sub>	L	VH	M	H	VH	H	M	H	H	H	H	M	L	VH	VH	M	H	H	H	H
C <sub>6</sub>	H	M	H	H	H	M	VH	H	H	H	VH	H	M	H	M	H	M	VH	M	H
C <sub>7</sub>	M	L	M	H	H	VL	H	H	M	L	VH	VH	M	M	H	H	VL	H	M	VH
C <sub>8</sub>	M	VL	M	H	H	H	L	H	M	VH	M	M	H	VH	H	H	H	M	VH	H
C <sub>9</sub>	H	VH	L	M	H	H	H	H	M	H	VH	H	M	VH	VH	H	VH	VH	H	VH
C <sub>10</sub>	M	M	VH	M	M	M	H	L	H	M	VH	L	M	VH	H	M	M	H	M	M
C <sub>11</sub>	M	VH	M	M	H	VH	H	H	H	VH	H	M	M	H	VH	M	H	VH	H	H
C <sub>12</sub>	H	VL	M	H	VH	VL	H	H	H	L	H	H	M	M	VH	H	M	M	H	H
C <sub>13</sub>	H	M	H	H	M	H	VH	M	VL	L	VH	H	M	M	L	H	M	M	M	M
C <sub>14</sub>	L	VH	M	VH	M	VH	H	VH	L	H	VH	H	M	M	M	VH	H	H	M	H
C <sub>15</sub>	M	H	M	VH	M	H	H	VH	H	M	H	VH	M	L	H	VH	M	VL	L	VH
C <sub>16</sub>	M	M	H	L	H	M	VH	L	M	H	VH	VH	M	H	M	H	M	L	M	M

Step 4: Apply fuzzy numbers (refer table -3)

Table 6: Fuzzy numbers for criteria rating

Criteria	DM <sub>1</sub>	DM <sub>2</sub>	DM <sub>3</sub>	DM <sub>4</sub>
C <sub>1</sub>	(5,7,9)	(7,9,9)	(7,9,9)	(5,7,9)
C <sub>2</sub>	(7,9,9)	(7,9,9)	(5,7,9)	(7,9,9)
C <sub>3</sub>	(7,9,9)	(7,9,9)	(5,7,9)	(7,9,9)
C <sub>4</sub>	(5,7,9)	(1,3,5)	(3,5,7)	(3,5,7)
C <sub>5</sub>	(5,7,9)	(7,9,9)	(7,9,9)	(7,9,9)
C <sub>6</sub>	(3,5,7)	(7,9,9)	(3,5,7)	(3,5,7)
C <sub>7</sub>	(3,5,7)	(1,1,3)	(3,5,7)	(1,3,5)
C <sub>8</sub>	(1,3,5)	(1,3,5)	(1,3,5)	(1,1,3)
C <sub>9</sub>	(5,7,9)	(5,7,9)	(3,5,7)	(3,5,7)
C <sub>10</sub>	(7,9,9)	(7,9,9)	(3,5,7)	(5,7,9)
C <sub>11</sub>	(5,7,9)	(5,7,9)	(5,7,9)	(5,7,9)
C <sub>12</sub>	(3,5,7)	(3,5,7)	(5,7,9)	(3,5,7)
C <sub>13</sub>	(5,7,9)	(1,3,5)	(3,5,7)	(3,5,7)
C <sub>14</sub>	(5,7,9)	(5,7,9)	(3,5,7)	(5,7,9)
C <sub>15</sub>	(3,5,7)	(5,7,9)	(3,5,7)	(3,5,7)
C <sub>16</sub>	(5,7,9)	(7,9,9)	(5,7,9)	(7,9,9)

Table 7: Fuzzy numbers for alternative rating

CRITERIA	S1				S2				S3				S4				S5			
	DM <sub>1</sub>	DM <sub>2</sub>	DM <sub>3</sub>	DM <sub>4</sub>	DM <sub>1</sub>	DM <sub>2</sub>	DM <sub>3</sub>	DM <sub>4</sub>	DM <sub>1</sub>	DM <sub>2</sub>	DM <sub>3</sub>	DM <sub>4</sub>	DM <sub>1</sub>	DM <sub>2</sub>	DM <sub>3</sub>	DM <sub>4</sub>	DM <sub>1</sub>	DM <sub>2</sub>	DM <sub>3</sub>	DM <sub>4</sub>
C <sub>1</sub>	(3,5,7)	(7,9,9)	(7,9,9)	(1,3,5)	(5,7,9)	(3,5,7)	(5,7,9)	(1,3,5)	(5,7,9)	(5,7,9)	(5,7,9)	(7,9,9)	(5,7,9)	(7,9,9)	(3,5,7)	(5,7,9)	(1,3,5)	(1,3,5)	(1,3,5)	(3,5,7)
C <sub>2</sub>	(5,7,9)	(7,9,9)	(5,7,9)	(1,3,5)	(5,7,9)	(7,9,9)	(5,7,9)	(1,3,5)	(5,7,9)	(5,7,9)	(7,9,9)	(5,7,9)	(5,7,9)	(5,7,9)	(3,5,7)	(5,7,9)	(5,7,9)	(3,5,7)	(3,5,7)	(3,5,7)
C <sub>3</sub>	(3,5,7)	(7,9,9)	(3,5,7)	(5,7,9)	(5,7,9)	(7,9,9)	(7,9,9)	(5,7,9)	(3,5,7)	(3,5,7)	(7,9,9)	(5,7,9)	(5,7,9)	(5,7,9)	(5,7,9)	(5,7,9)	(1,3,5)	(5,7,9)	(7,9,9)	(7,9,9)
C <sub>4</sub>	(1,1,3)	(1,1,3)	(5,7,9)	(5,7,9)	(3,5,7)	(1,3,5)	(7,9,9)	(5,7,9)	(3,5,7)	(5,7,9)	(7,9,9)	(5,7,9)	(3,5,7)	(3,5,7)	(5,7,9)	(7,9,9)	(3,5,7)	(7,9,9)	(1,3,5)	(7,9,9)
C <sub>5</sub>	(1,3,5)	(7,9,9)	(3,5,7)	(5,7,9)	(7,9,9)	(5,7,9)	(3,5,7)	(5,7,9)	(5,7,9)	(5,7,9)	(5,7,9)	(3,5,7)	(1,3,5)	(7,9,9)	(7,9,9)	(3,5,7)	(5,7,9)	(5,7,9)	(5,7,9)	(5,7,9)
C <sub>6</sub>	(5,7,9)	(3,5,7)	(5,7,9)	(5,7,9)	(5,7,9)	(3,5,7)	(7,9,9)	(5,7,9)	(5,7,9)	(5,7,9)	(7,9,9)	(5,7,9)	(3,5,7)	(5,7,9)	(3,5,7)	(5,7,9)	(3,5,7)	(7,9,9)	(3,5,7)	(5,7,9)
C <sub>7</sub>	(3,5,7)	(1,3,5)	(3,5,7)	(5,7,9)	(5,7,9)	(1,1,3)	(5,7,9)	(5,7,9)	(3,5,7)	(1,3,5)	(7,9,9)	(7,9,9)	(3,5,7)	(3,5,7)	(5,7,9)	(5,7,9)	(1,1,3)	(5,7,9)	(3,5,7)	(7,9,9)
C <sub>8</sub>	(3,5,7)	(1,1,3)	(3,5,7)	(5,7,9)	(5,7,9)	(5,7,9)	(1,3,5)	(5,7,9)	(3,5,7)	(7,9,9)	(3,5,7)	(5,7,9)	(5,7,9)	(7,9,9)	(5,7,9)	(5,7,9)	(5,7,9)	(3,5,7)	(7,9,9)	(5,7,9)
C <sub>9</sub>	(5,7,9)	(7,9,9)	(1,3,5)	(3,5,7)	(5,7,9)	(5,7,9)	(5,7,9)	(5,7,9)	(3,5,7)	(5,7,9)	(7,9,9)	(5,7,9)	(3,5,7)	(7,9,9)	(7,9,9)	(5,7,9)	(7,9,9)	(7,9,9)	(5,7,9)	(7,9,9)
C <sub>10</sub>	(3,5,7)	(3,5,7)	(7,9,9)	(3,5,7)	(3,5,7)	(3,5,7)	(5,7,9)	(1,3,5)	(5,7,9)	(3,5,7)	(7,9,9)	(1,3,5)	(3,5,7)	(7,9,9)	(5,7,9)	(3,5,7)	(3,5,7)	(5,7,9)	(3,5,7)	(3,5,7)
C <sub>11</sub>	(3,5,7)	(7,9,9)	(3,5,7)	(3,5,7)	(5,7,9)	(7,9,9)	(5,7,9)	(5,7,9)	(5,7,9)	(7,9,9)	(5,7,9)	(3,5,7)	(3,5,7)	(5,7,9)	(7,9,9)	(3,5,7)	(5,7,9)	(7,9,9)	(5,7,9)	(5,7,9)
C <sub>12</sub>	(5,7,9)	(1,1,3)	(3,5,7)	(5,7,9)	(7,9,9)	(1,1,3)	(5,7,9)	(5,7,9)	(5,7,9)	(1,3,5)	(5,7,9)	(5,7,9)	(3,5,7)	(3,5,7)	(7,9,9)	(5,7,9)	(3,5,7)	(3,5,7)	(5,7,9)	(5,7,9)
C <sub>13</sub>	(5,7,9)	(3,5,7)	(5,7,9)	(5,7,9)	(3,5,7)	(5,7,9)	(7,9,9)	(3,5,7)	(1,1,3)	(1,3,5)	(7,9,9)	(5,7,9)	(3,5,7)	(3,5,7)	(1,3,5)	(5,7,9)	(3,5,7)	(3,5,7)	(3,5,7)	(3,5,7)
C <sub>14</sub>	(1,3,5)	(7,9,9)	(3,5,7)	(7,9,9)	(3,5,7)	(7,9,9)	(5,7,9)	(7,9,9)	(1,3,5)	(5,7,9)	(7,9,9)	(5,7,9)	(3,5,7)	(3,5,7)	(3,5,7)	(7,9,9)	(5,7,9)	(5,7,9)	(3,5,7)	(5,7,9)
C <sub>15</sub>	(3,5,7)	(5,7,9)	(3,5,7)	(7,9,9)	(3,5,7)	(5,7,9)	(5,7,9)	(7,9,9)	(5,7,9)	(3,5,7)	(5,7,9)	(7,9,9)	(3,5,7)	(1,3,5)	(5,7,9)	(7,9,9)	(3,5,7)	(1,1,3)	(1,3,5)	(7,9,9)
C <sub>16</sub>	(3,5,7)	(3,5,7)	(7,9,9)	(1,3,5)	(5,7,9)	(3,5,7)	(7,9,9)	(1,3,5)	(3,5,7)	(5,7,9)	(7,9,9)	(7,9,9)	(3,5,7)	(5,7,9)	(3,5,7)	(5,7,9)	(3,5,7)	(1,3,5)	(3,5,7)	(3,5,7)

Step 5: Aggregated alternative and criteria weightage fuzzy decision matrix.

Table 8: Aggregated fuzzy decision matrix for alternative

Criteria	S <sub>1</sub>	S <sub>2</sub>	S <sub>3</sub>	S <sub>4</sub>	S <sub>5</sub>
C <sub>1</sub>	(1,6,5,9)	(1,5,5,9)	(5,7,5,9)	(3,7,9)	(1,3,5,7)
C <sub>2</sub>	(1,6,5,9)	(1,6,5,9)	(5,7,5,9)	(3,6,5,9)	(3,5,5,9)
C <sub>3</sub>	(3,6,5,9)	(5,8,9)	(3,6,5,9)	(5,7,9)	(1,6,5,9)
C <sub>4</sub>	(1,4,9)	(1,6,9)	(3,7,9)	(3,6,5,9)	(5,7,9)
C <sub>5</sub>	(1,6,9)	(3,7,9)	(3,7,5,9)	(1,6,5,9)	(3,6,5,9)
C <sub>6</sub>	(3,6,5,9)	(3,7,9)	(1,6,5,9)	(3,6,9)	(3,6,5,9)
C <sub>7</sub>	(1,5,9)	(1,5,5,9)	(1,6,5,9)	(3,6,9)	(1,5,5,9)
C <sub>8</sub>	(1,4,5,9)	(1,6,9)	(3,6,5,9)	(5,7,5,9)	(3,7,9)
C <sub>9</sub>	(1,6,9)	(5,7,9)	(3,7,5,9)	(3,7,5,9)	(5,7,9)
C <sub>10</sub>	(3,6,9)	(1,5,9)	(1,6,9)	(3,6,5,9)	(3,5,5,9)
C <sub>11</sub>	(3,6,9)	(5,7,5,9)	(3,7,9)	(3,6,5,9)	(5,7,5,9)
C <sub>12</sub>	(1,5,9)	(1,6,9)	(3,6,5,9)	(3,6,5,9)	(3,6,9)
C <sub>13</sub>	(3,6,5,9)	(3,6,5,9)	(1,5,9)	(1,5,9)	(3,5,7)
C <sub>14</sub>	(1,6,5,9)	(3,7,5,9)	(1,6,5,9)	(3,6,9)	(3,6,5,9)
C <sub>15</sub>	(3,6,5,9)	(3,7,9)	(3,7,9)	(1,6,9)	(1,4,5,9)
C <sub>16</sub>	(1,5,5,9)	(1,5,5,9)	(3,7,5,9)	(3,6,9)	(1,4,5,7)

Taking first criteria ( $C_1$ ) and first supplier value ( $S_1$ ) from table-7 (use equation 2)

1.  $a_{ij} = \min_k \{a_{kij}\} = 1.000$  [i.e. minimum value of first place (3,5,7 ; 7,9,9 ; 7,9,9 & 1,3,5)]
2.  $b_{ij} = 1/k \sum_{k=1}^k \{b_{kij}\} = 6.5$  [i.e. average value of middle place (3,5,7 ; 7,9,9 ; 7,9,9 & 1,3,5)]
3.  $c_{ij} = \max_k \{c_{kij}\} = 9.000$  [i.e. maximum value of last place (3,5,7 ; 7,9,9 ; 7,9,9 & 1,3,5)]

Same procedure has been used to evaluate table-9 by using table- 6 and equation 2.

**Table 9: Aggregated fuzzy decision matrix for criteria weightage**

Criteria	Aggregate Weightage
$C_1$	(5,8,9)
$C_2$	(5,8.5,9)
$C_3$	(5,8.5,9)
$C_4$	(1,5,9)
$C_5$	(5,8.5,9)
$C_6$	(3,6,9)
$C_7$	(1,3,5,7)

$C_8$	(1,2.5,5)
$C_9$	(3,4,9)
$C_{10}$	(3,7.5,9)
$C_{11}$	(5,7,9)
$C_{12}$	(3,5.5,9)
$C_{13}$	(1,5,9)
$C_{14}$	(3,6.5,9)
$C_{15}$	(3,5.5,9)
$C_{16}$	(5,8,9)

**Step 6:** Fuzzy multi criteria group decision making and process of normalizing.

As we are working on various criteria for decision-making, some might be benefit criteria and some might be cost criteria. Aim is to maximize benefit and minimize the cost. A fuzzy multi criteria group decision-making problem has been concisely expressed in matrix format using step 3 and step 4 of research methodology and also using equation 5 & 6.

In this study we considered criteria no. 4, 7, 8 and 14 under cost criteria and rest criteria 1, 2, 3, 5, 6, 9, 10, 11, 12, 13, 15 and 16 under benefit criteria.

**Table 10: Normalized aggregated fuzzy decision matrix for alternatives**

Criteria	$S_1$	$S_2$	$S_3$	$S_4$	$S_5$
$C_1$	(0.111,0.722,1)	(0.111,0.611,1)	(0.556,0.833,1)	(0.333,0.778,1)	(0.111,0.944,0.778)
$C_2$	(0.111,0.722,1)	(0.111,0.722,1)	(0.556,0.833,1)	(0.333,0.722,1)	(0.333,0.611,1)
$C_3$	(0.333,0.722,1)	(0.556,0.889,1)	(0.333,0.722,1)	(0.556,0.778,1)	(0.111,0.722,1)
$C_4$	(0.111,0.25,1)	(0.111,0.167,1)	(0.111,0.143,0.333)	(0.111,0.154,0.333)	(0.111,0.154,1)
$C_5$	(0.111,0.667,1)	(0.333,0.778,1)	(0.333,0.833,1)	(0.111,0.722,1)	(0.556,0.778,1)
$C_6$	(0.333,0.722,1)	(0.333,0.778,1)	(0.111,0.722,1)	(0.333,0.667,1)	(0.333,0.722,1)
$C_7$	(0.111,0.2,1)	(0.111,0.182,1)	(0.111,0.154,1)	(0.111,0.167,0.333)	(0.111,0.182,1)
$C_8$	(0.111,0.222,1)	(0.111,0.167,1)	(0.111,0.154,0.333)	(0.111,0.133,0.2)	(0.111,0.143,0.333)
$C_9$	(0.111,0.667,1)	(0.556,0.778,1)	(0.333,0.778,1)	(0.333,0.833,1)	(0.556,0.778,1)
$C_{10}$	(0.333,0.667,1)	(0.111,0.556,1)	(0.111,0.667,1)	(0.333,0.722,1)	(0.333,0.611,1)
$C_{11}$	(0.333,0.667,1)	(0.556,0.833,1)	(0.333,0.778,1)	(0.333,0.722,1)	(0.556,0.833,1)
$C_{12}$	(0.111,0.556,1)	(0.111,0.667,1)	(0.111,0.667,1)	(0.333,0.722,1)	(0.333,0.667,1)
$C_{13}$	(0.333,0.722,1)	(0.333,0.722,1)	(0.111,0.556,1)	(0.111,0.556,1)	(0.333,0.556,0.778)
$C_{14}$	(0.111,0.154,1)	(0.111,0.133,0.33)	(0.111,0.154,1)	(0.111,0.167,0.333)	(0.111,0.154,0.333)
$C_{15}$	(0.333,0.722,1)	(0.333,0.778,1)	(0.333,0.778,1)	(0.111,0.667,1)	(0.111,0.5,1)
$C_{16}$	(0.111,0.611,1)	(0.111,0.611,1)	(0.333,0.833,1)	(0.333,0.667,1)	(0.111,0.5,0.778)

The ranges of normalized triangular fuzzy numbers belong to (0,1).

Table 11: Weighted normalized fuzzy decision matrix.

Criteria	S <sub>1</sub>	S <sub>2</sub>	S <sub>3</sub>	S <sub>4</sub>	S <sub>5</sub>
C <sub>1</sub>	(0.556,5.776,9)	(0.556,4.889,9)	(2.78,6.664,9)	(1.665,6.224,9)	(0.555,7.552,7.002)
C <sub>2</sub>	(0.555,6.137,9)	(0.555,6.137,9)	(2.78,7.081,9)	(1.665,6.137,9)	(1.665,5.194,9)
C <sub>3</sub>	(1.665,6.137,9)	(2.787,0.557,9)	(1.665,6.137,9)	(2.78,6.613,9)	(0.555,6.137,9)
C <sub>4</sub>	(0.111,1.25,9)	(0.111,0.835,9)	(0.111,0.715,2.997)	(0.111,0.770,2.997)	(0.111,0.77,9)
C <sub>5</sub>	(0.555,5.670,9)	(1.665,6.613,9)	(1.665,7.081,9)	(0.555,,6.137,9)	(2.78,6.613,9)
C <sub>6</sub>	0.999,4.332,9	(0.999,4.668,9)	(0.333,4.332,9)	(0.999,4.002,9)	(0.999,4.332,9)
C <sub>7</sub>	0.111,0.7,7	(0.111,0.637,7)	(0.111,0.539,7)	(0.111,0.585,2.333)	(0.111,0.637,7)
C <sub>8</sub>	0.111,0.555,5	(0.111,0.418,5)	(0.111,0.385,1.665)	(0.111,0.333,1)	(0.111,0.358,1.665)
C <sub>9</sub>	0.333,2.668,9	(1.668,3.112,9)	(0.999,3.112,9)	(0.999,3.332,9)	(1.668,3.112,9)
C <sub>10</sub>	0.999,5.003,9	(0.333,4.17,9)	(0.333,5.003,9)	(0.999,5.415,9)	(0.999,4.583,9)
C <sub>11</sub>	1.665,4.669,9	(2.78,5.831,9)	(1.665,5.446,9)	(1.665,5.054,9)	(2.78,5.831,9)
C <sub>12</sub>	0.333,3.058,9	(0.333,3.669,9)	(0.333,3.669,9)	(0.999,3.971,9)	(0.999,3.669,9)
C <sub>13</sub>	0.333,3.61,9	(0.333,3.61,9)	(0.111,2.78,9)	(0.555,2.78,9)	(0.333,2.78,7.002)
C <sub>14</sub>	(0.333,4.693,9)	(0.333,0.865,2.997)	(0.999,5.057,9)	(0.333,4.336,9)	(0.333,1.001,2.997)
C <sub>15</sub>	(0.999,3.971,9)	(0.999,4.279,9)	(0.999,4.279,9)	(0.333,3.669,9)	(0.333,2.75,9)
C <sub>16</sub>	(0.555,4.888,9)	(0.555,4.888,9)	(1.665,6.664,9)	(1.665,5.336,9)	(0.555,4,7.002)

Table-11 has been evaluated by using data of table-9, table-10 and applying it in equation 7.

#### Step 7: FPIS and FNIS

Using equation number 8 & 9, we have selected the maximum value from each row as  $p^+$  and minimum value from each row as  $p^-$  as shown in table-11. Table-12 shows both these values.

Table 12: Final list of values of  $p^+$  and  $p^-$  for each alternative

A <sup>+</sup>	A <sup>-</sup>
P <sub>1</sub> <sup>+</sup> (9,9,9)	P <sub>1</sub> <sup>-</sup> (0.555,0.555,0.555)
P <sub>2</sub> <sup>+</sup> (9,9,9)	P <sub>2</sub> <sup>-</sup> (0.555,0.555,0.555)
P <sub>3</sub> <sup>+</sup> (9,9,9)	P <sub>3</sub> <sup>-</sup> (0.555,0.555,0.555)
P <sub>4</sub> <sup>+</sup> (9,9,9)	P <sub>4</sub> <sup>-</sup> (0.111,0.111,0.111)
P <sub>5</sub> <sup>+</sup> (9,9,9)	P <sub>5</sub> <sup>-</sup> (0.555,0.555,0.555)
P <sub>6</sub> <sup>+</sup> (9,9,9)	P <sub>6</sub> <sup>-</sup> (0.333,0.333,0.333)
P <sub>7</sub> <sup>+</sup> (7,7,7)	P <sub>7</sub> <sup>-</sup> (0.111,0.111,0.111)
P <sub>8</sub> <sup>+</sup> (5,5,5)	P <sub>8</sub> <sup>-</sup> (0.111,0.111,0.111)
P <sub>9</sub> <sup>+</sup> (9,9,9)	P <sub>9</sub> <sup>-</sup> (0.333,0.333,0.333)

P <sub>10</sub> <sup>+</sup> (9,9,9)	P <sub>10</sub> <sup>-</sup> (0.333,0.333,0.333)
P <sub>11</sub> <sup>+</sup> (9,9,9)	P <sub>11</sub> <sup>-</sup> (1.665,1.665,1.665)
P <sub>12</sub> <sup>+</sup> (9,9,9)	P <sub>12</sub> <sup>-</sup> (0.333,0.333,0.333)
P <sub>13</sub> <sup>+</sup> (9,9,9)	P <sub>13</sub> <sup>-</sup> (0.111,0.111,0.111)
P <sub>14</sub> <sup>+</sup> (9,9,9)	P <sub>14</sub> <sup>-</sup> (0.333,0.333,0.333)
P <sub>15</sub> <sup>+</sup> (9,9,9)	P <sub>15</sub> <sup>-</sup> (0.333,0.333,0.333)
P <sub>16</sub> <sup>+</sup> (9,9,9)	P <sub>16</sub> <sup>-</sup> (0.555,0.555,0.555)

#### Step 8: FPIS and FNIS for each criteria

Here, we have to use equation-1 and table-12. Then, we have to find distance of each criterion from FPIS and FNIS for five alternatives.

Table 13: Distance of criteria of each alternative from FPIS and FNIS

Criteria	FPIS(S <sub>1</sub> )	FNIS(S <sub>1</sub> )	FPIS(S <sub>2</sub> )	FNIS(S <sub>2</sub> )	FPIS(S <sub>3</sub> )	FNIS(S <sub>3</sub> )	FPIS(S <sub>4</sub> )	FNIS(S <sub>4</sub> )	FPIS(S <sub>5</sub> )	FNIS(S <sub>5</sub> )
C <sub>1</sub>	5.218	5.732	5.422	5.480	3.836	6.153	4.528	5.907	5.079	5.493
C <sub>2</sub>	5.148	5.845	5.148	5.845	3.758	6.294	4.546	5.879	4.771	5.600
C <sub>3</sub>	4.546	5.879	3.686	6.463	4.546	4.883	3.846	6.136	5.148	5.845
C <sub>4</sub>	6.808	5.174	6.969	5.145	7.825	1.702	7.805	1.709	6.994	5.146
C <sub>5</sub>	5.241	5.700	4.453	6.035	4.377	6.195	5.148	5.845	3.846	6.136
C <sub>6</sub>	5.348	5.524	5.253	5.608	5.684	5.511	5.447	5.447	5.348	5.524
C <sub>7</sub>	5.389	3.992	5.414	3.989	5.453	3.985	6.067	1.311	5.414	3.989
C <sub>8</sub>	3.815	2.834	3.869	2.828	3.992	3.985	4.534	.530	4.343	0.908
C <sub>9</sub>	6.198	5.184	5.429	5.311	5.735	5.269	5.661	5.309	5.429	5.311
C <sub>10</sub>	5.164	5.697	5.728	5.472	5.510	5.684	5.062	5.813	5.276	5.586
C <sub>11</sub>	4.917	4.576	4.030	4.913	4.706	4.764	4.809	4.665	4.030	4.913
C <sub>12</sub>	6.067	5.245	5.875	5.362	5.875	5.362	5.456	5.440	5.551	5.375
C <sub>13</sub>	5.893	5.516	5.893	5.517	6.264	5.358	6.055	5.365	6.266	4.268
C <sub>14</sub>	5.588	5.601	7.688	1.568	5.150	5.712	5.682	5.512	7.641	1.585
C <sub>15</sub>	5.456	5.440	5.364	5.512	5.364	5.512	5.875	5.362	6.169	5.195
C <sub>16</sub>	5.423	5.480	5.423	5.480	4.444	6.052	4.734	5.640	7.309	4.220

For evaluating first cell value of table-13, we consider first cell C<sub>1</sub> of table-11 and then calculating distance of criteria for that alternative.

Criteria	S <sub>1</sub>
C <sub>1</sub>	(0.556,5.776,9)

From table-12,  $p_1^+(9, 9, 9)$

From equation-1

Now, for (C<sub>1</sub>-S<sub>1</sub>) from table-11 and P<sub>1</sub><sup>+</sup> from table-12

$$d = \sqrt{1/3 [(0.556 - 9)^2 + (5.776 - 9)^2 + (9 - 9)^2]}$$

= 5.218 (likewise use same process to find remaining values of table-13)

**Step 9:** The distance of each weighted alternative (by using equation 10 and 11)

$$d_1^+ = 86.219, d_2^+ = 85.644, d_3^+ = 82.518, d_4^+ = 85.255, d_5^+ = 88.614$$

$$d_1^- = 83.419, d_2^- = 80.528, d_3^- = 82.421, d_4^- = 75.87, d_5^- = 75.094$$

**Step 10:** Closeness coefficient of each alternative (by using equation 12)

$$CC_1 = 0.492, CC_2 = 0.485, CC_3 = 0.50, CC_4 = 0.471, CC_5 = 0.459$$

## RESULTS AND DISCUSSION

In this study fuzzy TOPSIS technique has been adopted. Study has been conducted with five alternatives S<sub>1</sub>, S<sub>2</sub>, S<sub>3</sub>, S<sub>4</sub> and S<sub>5</sub>

with sixteen criteria C<sub>1</sub>, C<sub>2</sub>, C<sub>3</sub>, ..... and C<sub>16</sub> and four decision makers namely DM<sub>1</sub>, DM<sub>2</sub>, DM<sub>3</sub> and DM<sub>4</sub>. Different suppliers have been ranked according to the closeness coefficient CC<sub>i</sub> in decreasing order. The best alternative is closest to the FPIS and farthest from the FNIS. The ranking order of S<sub>1</sub>, S<sub>2</sub>, S<sub>3</sub>, S<sub>4</sub> and S<sub>5</sub> has been tabulated in table-14.

Table: 14 Rank assigned to suppliers

Alternatives	CC <sub>i</sub>	Rank
Supplier <sub>1</sub>	0.492	2
Supplier <sub>2</sub>	0.485	3
Supplier <sub>3</sub>	0.50	1
Supplier <sub>4</sub>	0.471	4
Supplier <sub>5</sub>	0.459	5

It was found that the supplier S<sub>3</sub> to be the best supplier among the suppliers. It doesn't justify that S<sub>1</sub>, S<sub>2</sub>, S<sub>4</sub> and S<sub>5</sub> are the efficient. The results may vary, based on the criterion priority levels by decision makers as analysis was carried out based on the criteria suggested by the various decision makers. The final results have been shown in table-14. By comparing CC<sub>i</sub> values of the five alternatives, we find that S<sub>3</sub> > S<sub>1</sub> > S<sub>2</sub> > S<sub>4</sub> > S<sub>5</sub>. S<sub>3</sub> is the Best choice considering the given criteria and S<sub>5</sub> is the worst choice considering given criteria. The idea for decisions about remaining suppliers S<sub>1</sub>, S<sub>2</sub>, and S<sub>4</sub> can be obtained from the rank assigned to them in table-14.

## CONCLUSION:

This paper discusses a study with an objective to analyze, how

to select the most efficient supplier in Indian pharmaceutical industry by fuzzy TOPSIS method in supplier MCDM problems when decision makers set the target value of each criterion. In this study, total 16 criteria were considered and a multi-criteria decision-making approach has been used for the supplier rating under fuzzy environment. This study approach comprises of two steps. In step first, the criteria for suppliers rating selection are identified. In second step the experts in pharmaceutical industry were asked to give linguistic ratings to the criteria and the alternatives. This study proposed a method and a procedure to extend the TOPSIS method to solve the problem. The results obtained from this case study may be used for decision-making of supplier selection rating in Indian pharmaceutical industry. The suppliers rating problem discussed in this paper can also be solved by Fuzzy AHP and the obtained results can be compared with the results obtained using Fuzzy TOPSIS method.

## REFERENCES

- [1] Athena Forghani, S. j. (2018), *A supplier selection in pharmaceutical supply chain using PCA, Z-TOPSIS and MILP: A case study, public library of science (PLOS) One*, 1-17.
- [2] Chris I. E., Bell-Hanyes J. (2010), *A model for quantifying strategic supplier selection: evidence from a generic pharmaceutical firm supply chain, international journal of business, marketing, and decision sciences*, 45-57.
- [3] David Asamoah, J. A. (2012), *AHP Approach for Supplier Evaluation and Selection in Pharmaceutical Manufacturing firm in Ghana, International Journal of Business and Management*, 49-62.
- [4] Enyinda, C. I., Dunu, E., & Gebremikael, F. (2010), *An analysis of strategic supplier selection and evaluation in a generic pharmaceutical firm supply chain, ASBBS Annual Conference: Las Vegas*, 77-91.
- [5] Farimah mokhatab rafiei, N. M. (2012), *Application of fuzzy TOPSIS method for the selection of warehouse location, A Case Study, Interdisciplinary Journal of Contemporary Research in Business*, 655-671.
- [6] J.M. Benitez, J.C. Martin, and C. Roman (2007), *Using fuzzy number for measuring quality of service in the hotel industry, Tourism Management*, 544-555.
- [7] Kore, M. N. (2017), *A Simplified Description of Fuzzy TOPSIS Method for Multi Criteria Decision Making, International Research Journal of Engineering and Technology (IRJET)*, 56-72.
- [8] Liao C-N, Kao H-P (2011), *An integrated fuzzy TOPSIS and MCGP approach to supplier selection in supply chain management, Expert Syst Appl*, 38-51.
- [9] M. Yurdakul, and Y.T. Ic (2008), *Analysis of the benefit generated by using fuzzy numbers in a TOPSIS model developed for machine tool selection problems, Journal of Materials Processing Technology*, 324-339.
- [10] Mehmet Sevkli, S. Z. (2014), *An application of fuzzy TOPSIS method for supplier selection, IEEE International Conference Fuzzy systems*, 978-985.
- [11] P. Kelle, J. Woosley, and H. Schneider (2012), *Pharmaceutical supply chain specifics and inventory solutions for a hospital case, Oper. Res. Heal. Care*, vol. 54-63.
- [12] Rouyendegh BD, Saputro TE. (2014), *Supplier Selection Using Integrated Fuzzy TOPSIS and MCGP, A Case Study, Procd Soc Behv.*, 3957-4027.
- [13] S. Onut., and S. Soner (2008), *Transshipment site selection using the AHP and TOPSIS approaches under fuzzy environment, Waste Management*, 1552-1559.
- [14] Tahriri, F., Osman, M. R., Ali, A., & Mohd, R (2008), *A review of supplier selection methods in manufacturing industries, Suranaree Journal of Science and Technology*, 201-208.
- [15] T.C. Chu, and Y.C. Lin (2003), *A fuzzy TOPSIS method for robot selection, The International Journal of Advanced Manufacturing Technology*, 284-290.
- [16] T.C. Wang, and T.H. Chang (2007), *Application of TOPSIS in evaluating initial training aircraft under fuzzy environment, Expert Systems with Applications*, 870-880.
- [17] T.C. Chu (2002), *Selecting plant location via a fuzzy TOPSIS approach, International Journal of Advanced Manufacturing Technology*, 859-864.
- [18] T. Yang, and C.C. Hung (2007), *Multiple-attribute decision making methods for plant layout design problem, Robotics and Computer-Integrated Manufacturing*, 126-137.
- [19] tonekaboni, k. s. (2012), *Application of topsis method to supplier selection in iran auto supply chain, Journal of Global Strategic Management*, 123-131.
- [20] Vinod Yadav, M. K. (2016), *Intelligent evaluation of suppliers using extent fuzzy TOPSIS method, A case study of an indian manufacturing SME, emerald publishing limited*, 0114-0143.
- [21] Weber C.A., Current J.R., Benton W.C. (1991), *Supplier selection criteria and methods, European Journal of Operational Research*, 2-18.

## AUTHORS

**Dashrath Kumar Patidar**, Research Scholar, Ujjain Engineering College, Ujjain - Indore Road, Vasant Vihar, Ujjain – 456 010, (MP)  
Email: patidardashrath42@gmail.com

**Dr. V. K. Sukhwani**, Professor, Department of Mechanical Engineering, Ujjain Engineering College, Ujjain - Indore Road, Vasant Vihar, Ujjain – 456 010, (MP)  
Email: v1sukhwani@gmail.com

**Dr. A. C. Shukla**, Professor, Department of Mechanical Engineering, Ujjain Engineering College, Ujjain - Indore Road, Vasant Vihar, Ujjain – 456 010, (MP)  
Email: apratuls@gmail.com